ANIL AGGRAWAL'S
Internet Journal of Forensic Medicine and Toxicology
The world's first online journal devoted entirely to forensic medicine and toxicology · Published 1 January and 1 July

VOL. 3, NO. 2
Downloaded from www.anilaggrawal.com · ISSN 0972-8074 (Online) · https://www.anilaggrawal.com/ij/vol_003_no_002/others/thesis_avneesh/chap005_2.html

Sex distribution

Chapters at a glance
[chapter 1] [chapter 2] [chapter 3] [chapter 4]

chapter 5 [Part 1] [Part 2] [Part 3] [Part 4] [Part 5] [Part 6]

[chapter 6] [chapter 7] [chapter 8] [chapter 9] [Photographs]

Back to the journal

SEX DISTRIBUTION

Males comprised a majority and constituted 98 (94.2%) compared to females who were only 6 (5.7%). The male to female ratio in the study was 16.3:1 (Male=98, Female=6). Male to female ratio in the most commonly affected age group 31-40 years was 13.5:1.

Age range	Male	Female
0-10	5	-
11-20	15	1
21-30	22	1
31-40	22	2
41-50	18	1
51-60	10	-
61-70	4	1
71-80	2	-
Total	98	6

TIME OF INCIDENCE

Most of the incidents irrespective of the cause, occurred between 9PM to 12 mid night comprising 28.8% of total cases and a large percentage of cases 41% occurred between 6PM to 12 mid night.

The least number of incidents occurred between 3AM to 6AM comprising 3% of the total cases. While in nine cases the time of incidence was not known.

Time of Incidence	No. of victims	%
12:01 – 03:00 AM	12	11.5
03:01 – 06:00 AM	4	3.8
06:01 – 09:00 AM	10	9.6
09:01 – 12:00 Noon	2	1.9
12:01 – 03:00 PM	5	4.8
03:01 – 06:00 PM	9	8.6
06:01 – 09:00 PM	13	12.5
09:01 – 12:00 PM	30	28.8
Unknown time	9	8.6